

Oculomotor Workshop

Timothy C. Hain, MD

Video Frenzel Goggles



- Expensive (about \$2000)
- Good teaching tool
- Can do some things not easily done with optical system (i.e vibration test, hyperventilation, vertebral artery test, head prone test, cross-cover)

Dr Hain's system



3

Optical Frenzel Goggles



- Inexpensive (about \$500)
- Portable – take to hospital
- A little limited – can't do vibration, head-forward or cross-cover
- Can get hot, bulbs burn out and break

4

Video Eye Movement Tests

- Saccades (slow, omn palsy, dysmetric)
- Spontaneous nystagmus
- Vibration
- Head-shaking
- Gaze testing
- Positional testing


5

Slow saccades



6


Case



Source: Leigh/Zee

7


Case



Source: Leigh/Zee

8


Case



Source: Leigh/Zee

9

Case




Source of video: Preston Calvert, M.D.

10

INO (Internuclear ophthalmoplegia)

- Brainstem lesion of MLF
- Most commonly seen in MS
- Slowing of adducting saccades
- Overshoot of abducting eye
- A system that can visualize both eyes is best



11

Oculomotor palsies

- Operators of ENG equipment may not think to do an oculomotor exam or check visual acuity.
- Tests that displace eye into paretic field may produce very asymmetrical results.

12

Case


Saccadic Dysmetria



13

Overshoot dysmetria

- Usually cerebellar lesion
- Occasionally parietic eye fixation
- Never peripheral vestibular lesion



14

Case

- 14 year old girl
- Very unstable gait
- headaches
- Darting eyes




15

Opsoclonus

- “dancing eyes-dancing feet” pediatric syndrome
- Neuroblastoma
- Paraneoplastic syndrome
- West Nile

16

CASE




17

Case




18

Cases



19

Case



20

Spontaneous Nystagmus

- Acute vestibular disorders (V Neuritis, horizontal canal BPPV, Menieres, recent surgery) have strong horizontal “jerk” nystagmus.
- Normal people and chronic vestibular disorders have little or no nystagmus. Neural compensation for vestibular tone asymmetry is fast and effective.
- Most people can't “fake” nystagmus.
- Almost everything unusual is central.

21

Case – cross-cover test




22

Non-vestibular spontaneous nystagmus the common variants Latent Nystagmus

- Found in persons with congenital esotropia
- changes direction according to viewing eye (Cross-cover test)
- Viewing eye beats laterally
- Intent to view controls direction (pseudoscope)
- Always have “lazy” eye

23

Latent Nystagmus



Note “bizzare” increasing velocity waveform typical of CN.
Some malingers use LN

24

Case



In light



In dark

25

Congenital Nystagmus

- One/1000 population
- Present from early age
- Usually worse in light
- PT not useful
- Rehab significance is to avoid confusing it with central nystagmus or vestibular nystagmus.

26

Non-vestibular spontaneous nystagmus: the common variants

- “Wrongly” directed primary position nystagmus
 - Downbeat
 - Upbeat
 - Torsional

27

Case

- Chiari (MRI)
- Cerebellar (especially remote effect) – get a CXR
- Idiopathic/drug




28

Case



29

The cause



30

Case

- Smoking (slight)
- Paxil (slight)
- Wernickes
- BPPV variants ?
- Vestibular neuritis variants
- Central vertigo – Migraine ?



31

Vibration test



Vibration test

- Method: Apply 60-120 hz vibration to SCM, first one side, then the other. Shower massagers work well for this and are inexpensive.
- Video Frenzel goggles – optical Frenzels don't work very well
- Compare nystagmus before and during



33

Vibration Induced Nystagmus

Vibration Induced Nystagmus in Unilateral Vestibular Loss

(c) 2005 Timothy C. Hain, M.D.

34

Vibration Induced Nystagmus

- Unidirectional horizontal nystagmus strongly suggests contralateral vestibular lesion.
- Permanent nystagmus – never goes away
- Direction changing nystagmus is a normal variant.
- Vertical or torsional nystagmus is of uncertain meaning. Seems more common in BPPV.

35


Head-shaking test

- Method: 20 cycles of horizontal head rotation
- Frenzel goggles to monitor nystagmus prior to and following head-shaking.
- Positive – substantial change in nystagmus following head-shaking. Usually beats away from bad ear.



36

Head-shaking in person with left sided vestibulopathy




37

HSN – unilat comments

- SN, HSN and Vibration are all useful in detecting Unilateral Vestibular Loss
- SN is seen acutely but vanishes over time.
- HSN is more sensitive to moderate loss than VN. However, it may appear and then vanish, or even go in wrong direction.
- Vibration is more dependable than HSN – never goes away.

38

Head Shaking in Bilateral Loss



39

Head Shaking in Bilateral Loss Suppression test



40

Head Shaking

- Moderately useful test –
- About 75% localizing
- Absent in about 25%
- No nystagmus – good for bilaterals

41

Gaze Testing

- Move finger to the limits of lateral gaze (bury sclera) – if can't bury, may have oculomotor palsy
- Move finger to limits of vertical gaze
- Do eyes reach end-gaze ?
- Is there end-gaze nystagmus ?
- Is there rebound nystagmus ?

42

Gaze Test: normal

- Minimal or no horizontal and upgaze nystagmus
- No down-gaze nystagmus in normal people
- No rebound nystagmus

43

Case (Cerebellar patient)

Patient BM
12/2003
Cerebellar
Degeneration

44

Rebound Nystagmus

- Nearly always cerebellar lesion
- Rarely congenital
- Method of separating out cerebellar GEN from sedative effect or congenital nystagmus

45

Positional Testing Strategies

- Dix-Hallpike --- head 30 back/rotated
 - Posterior canal BPPV (UBN, ipsitorision)
 - Anterior canal BPPV (DBN)
- Head 30deg fwd or supine – lateral canal BPPV
 - Geotropic or ageotropic
- Head upright or forward – cervical vertigo
 - Gravity coordinate vs. not

46

A Mat Table Helps



- Stable and safe
- Big enough to roll patients
- Locate emesis basin before beginning



47

Posterior Canal BPPV



48

Posterior Canal BPPV

- Upbeating/Torsional nystagmus (or at least torsional, top of eye beats toward ground)
- Latency: 0 to 30 sec
- Burst: up to 1 min
- Unwinds when sit up
- Treat with Epley/Semont/Brandt-Daroff

49

Direction Changing Positional Nystagmus



50

DCPN – Lateral Canal BPPV ?

- Geotropic vs Ageotropic
- Usually prolonged
- Reverses sense with head forward (cervical vertigo doesn't reverse)
- Treat with log-roll

51

Conclusion

- Video goggles are very useful !
- Removes fixation
- Large image
- Vibration test and head-forward test possible
- Impress patient and family

52

More details

Hain, T.C. Approach to the patient with Dizziness and Vertigo. Practical Neurology (Ed. Biller), 2002. Lippincott-Raven

More movies

www.dizziness-and-balance.com